arXiv:1005.0585v2 [math.DS] 20 Apr 2011

Minimal C^1-diffeomorphisms of the circle which admit measurable fundamental domains

Hiroki Kodama and Shigenori Matsumoto

Abstract. We construct, for each irrational number α, a minimal C^1-diffeomorphism of the circle with rotation number α which admits a measurable fundamental domain with respect to the Lebesgue measure.

1. Introduction

The concept of ergodicity is important not only for measure preserving dynamical systems but also for systems which admits a natural quasi-invariant measure. Given a probability space (X, μ) and a transformation T of X, μ is said to be *quasi-invariant* if the push forward $T_*\mu$ is equivalent to μ. In this case T is called *ergodic* with respect to μ, if a T-invariant Borel subset in X is either null or conull.

A diffeomorphism of a differentiable manifold always leaves the Riemannian volume (also called the Lebesgue measure) quasi-invariant, and one can ask if a given diffeomorphism is ergodic with respect to the Lebesgue measure (below *ergodic* for short) or not. Answering a question of A. Denjoy [**D**], M. Herman (Chapt. VII, p. 86, [**H**]) showed that a C^1-diffeomorphism of the circle with derivative of bounded variation is ergodic provided its rotation number is irrational. See also Chapt. 12.7, p. 419, [**KH**]. Contrarily Oliveira and da Rocha [**OR**] constructed a minimal C^1 diffeomorphism of the circle which is not ergodic.

At the opposite extreme of the ergodicity lies the concept of measurable fundamental domains. Given a transformation T of a standard probability space (X, μ) leaving μ quasi-invariant, a Borel subset C of X is called a *measurable fundamental domain* if T^nC $(n \in \mathbb{Z})$ is mutually disjoint and the union $\cup_{n\in\mathbb{Z}} T^nC$ is conull. In this case any Borel function on C can be extended to a T-invariant measurable function on X, and an ergodic component of T is just a single orbit. The purpose of this paper is to show the following theorem.

2000 *Mathematics Subject Classification.* Primary 37E15, secondary 37C05, 37A40.

Key words and phrases. diffeomorphism, minimality, rotation number, ergodicity.

The first author is partially supported by the Japan Society for the Promotion of Science (JSPS) through its "Funding Program for World-Leading Innovative R&D on Science and Technology (FIRST Program)." The second author is partially supported by Grant-in-Aid for Scientific Research (C) No. 20540096.

Theorem 1.1. *For any irrational number α, there is a minimal C^1-diffeomorphism of the circle with rotation number α which admits a measurable fundamental domain with respect to the Lebesgue measure.*

The rest of the paper is devoted to the proof of Theorem 1.1.

2. A measurable fundamental domain for a Lipschitz homeomorphism

For the sake of simplicity, we first consider the case of $\alpha = (\sqrt{5}-1)/2$, the inverse of golden ratio. We regard the circle S^1 as $\mathbb{R}/\mathbb{Z}$. Suppose R denotes the rotation by α. Define a Cantor set C in the circle by

$$C = \Big\{ \sum_{k=1}^{\infty} \frac{\varepsilon_k}{2^{3^k}} \ \Big|\ \varepsilon_k = 0 \text{ or } 1 \Big\} \pmod{\mathbb{Z}}.$$

Note that all numbers in C are well approximable by rational numbers.

Claim 2.1. *$R^nC \cap R^mC = \emptyset$ for any integers $n \neq m$.*

Proof. Suppose $x \in R^nC \cap R^mC$, then $x - n\alpha, x - m\alpha \in C$, therefore

$$(-n+m)\alpha \in C + (-C) := \Big\{ \sum_{k\geq 3} \frac{\varepsilon'_k}{2^{3^k}} \ \Big|\ \varepsilon'_k = 0 \text{ or } \pm 1 \Big\} \pmod{\mathbb{Z}}.$$

$(-n+m)\alpha$ is badly approximable, while $C + (-C)$ consists of well approximable numbers, which is a contradiction. □

See Section 4 for the detail and the general case.

Fix a probability measure μ_0 on C without atom such that $\mathrm{supp}(\mu_0) = C$. We also choose a sequence $(a_i)_{i\in\mathbb{Z}}$ of positive numbers satisfying $\sum_{i\in\mathbb{Z}} a_i = 1$. Now we can define a probability measure μ on S^1 by

$$\mu := \sum_{i\in\mathbb{Z}} a_i R^i_* \mu_0.$$

The Radon-Nikodym derivative $\frac{dR_*^{-1}\mu}{d\mu}$ is equal to $\frac{a_{i+1}}{a_i}$ on the set R^iC. Now we assume that $\frac{a_{i+1}}{a_i} \in [\frac{1}{D}, D]$ for some $D > 1$, then it follows that $\frac{dR_*^{-1}\mu}{d\mu} \in L^\infty(S^1, \mu)$.

We define a homeomorphism h of S^1 by $h(0) = 0$ and $h(x) = y$ if and only if $Leb[0,x] = \mu[0,y]$, where Leb denotes the Lebesgue measure on S^1; or more briefly, $h_* Leb = \mu$. Finally define a homeomorphism F of S^1 by $F := h^{-1} \circ R \circ h$, then

$$\frac{dF_*^{-1}\, Leb}{d\, Leb} = \frac{dR_*^{-1}\mu}{d\mu} \circ h \in L^\infty(S^1, Leb),$$

i.e. the map F is a Lipschitz homeomorphism. The set $C' = h^{-1}C$ is a measurable fundamental domain of F.

3. Make it C^1

3.1. What shall we do? To prove the Theorem, we are trying to make the Radon-Nikodym derivative $g = \frac{dR_*^{-1}\mu}{d\mu}$ continuous on S^1.

Fix an arbitrary point $x_0 \in C$. For a positive integer i,

$$\begin{aligned} a_i &= (a_i/a_{i-1})\cdots(a_2/a_1)(a_1/a_0)a_0 \\ &= g(R^{i-1}x_0)\cdots g(Rx_0)g(x_0)a_0, \\ a_{-i} &= (a_{-i+1}/a_{-i})^{-1}\cdots(a_{-1}/a_{-2})^{-1}(a_0/a_{-1})^{-1}a_0 \\ &= g(R^{-i}x_0)^{-1}\cdots g(R^{-2}x_0)^{-1}g(R^{-1}x_0)^{-1}a_0. \end{aligned}$$

Set $\phi = \log g$ and define a map $\Phi\colon S^1\times\mathbb{R}\to S^1\times\mathbb{R}$ by $\Phi(x,y) = (Rx, y+\phi(x))$. Simple calculation shows that $\Phi^n(x,y) = (R^n x, y+\phi^{(n)}(x))$ where

$$\begin{aligned} \phi^{(m)}(x) &= \sum_{i=0}^{m-1}\phi(R^i x) \qquad (m>0), \\ \phi^{(-m)}(x) &= -\sum_{i=1}^{m}\phi(R^{-i}x) \qquad (m>0), \\ \phi^{(0)}(x) &= 0. \end{aligned} \tag{3.1}$$

Therefore $a_i = \exp(\phi^{(i)}(x_0))a_0$. To satisfy $\sum_{i\in\mathbb{Z}} a_i = 1$, it suffice to find ϕ so that $\sum_{i\in\mathbb{Z}}\exp(\phi^{(i)}(x_0)) < \infty$.

3.2. Construction step I. Now we forget about a_i's and the Cantor set C. As a first step, we are going to construct a function $\phi\in C(S^1)$ satisfying $\sum_{i\in\mathbb{Z}}\exp(\phi^{(i)}(x_0)) < \infty$ for a single point x_0, where $\phi^{(i)}$ are defined by (3.1).

We are going to define continuous functions $\phi_n \in C(S^1)$ $(n\in\mathbb{N})$ in such a way that $\sum_{i=1}^{\infty}\|\phi_n\|_\infty < \infty$. Then $\phi = \sum_{i=1}^{\infty}\phi_n$ converges uniformly, thus ϕ is also continuous.

Fix an integer $n\in\mathbb{N}$. Choose a sufficiently small neighbourhood J of x_0 so that $R^{-2^n}J,\ldots,R^{-1}J,J,RJ,\ldots,R^{2^n-1}J$ are disjoint. Consider a bump function f on J so that $\operatorname{supp} f\subset J$, $f(x_0) = (3/4)^n$ and $0\le f(x) < (3/4)^n$ on $J\setminus\{x_0\}$. Define $\phi_n\colon S^1\to\mathbb{R}$ by

$$\phi_n(x) = \begin{cases} -f(R^{-i}x) & x\in R^iJ,\ i=0,1,\ldots,2^n-1 \\ f(R^{-i}x) & x\in R^iJ,\ i=-2^n,-2^n+1,\ldots,-1 \\ 0 & \text{otherwise.} \end{cases}$$

Lemma 3.1. $\phi_n^{(i)}(x_0)\begin{cases} = -|i|(3/4)^n & \textit{for } -2^n\le i\le 2^n \\ \le 0 & \textit{for any } i\in\mathbb{Z}. \end{cases}$

Proof. The equality for the first case is trivial. Define an increasing sequence $(m_k)_{k\in\mathbb{Z}}$ by $m_0 = 0$ and $\{m_k | k\in\mathbb{Z}\} = \{m\in\mathbb{Z} | R^m x_0\in J\}$. Since $R^{-2^n}J,\ldots,R^{2^n-1}J$ are disjoint, $m_{k+1}-m_k\ge 2^{n+1}$ for any $k\in\mathbb{Z}$. Using this sequence, $R^m x_0\in R^iJ$ if and only if $m = m_k + i$ for some k. Therefore,

$$\phi^{(i+1)}(x_0) = \begin{cases} \phi^{(i)}(x_0) & m_{k-1}+2^n\le i< m_k-2^n \\ \phi^{(i)}(x_0) + f(R^{m_k}x_0) & m_k-2^n\le i< m_k \\ \phi^{(i)}(x_0) - f(R^{m_k}x_0) & m_k\le i< m_k+2^n \end{cases}$$



FIGURE 1. The graph of ϕ_n and the orbit of x_0

for some k. Induction for $|k|$ shows that

$$\phi^{(i)}(x_0) = \begin{cases} -2^n \cdot (3/4)^n & m_{k-1} + 2^n \leq i \leq m_k - 2^n \\ -2^n \cdot (3/4)^n + (i - (m_k - 2^n))f(R^{m_k}x_0) & m_k - 2^n \leq i \leq m_k \\ -2^n \cdot (3/4)^n + ((m_k + 2^n) - i)f(R^{m_k}x_0) & m_k \leq i \leq m_k + 2^n. \end{cases}$$

Since $f(R^{m_k}x_0) \leq (3/4)^n$, the inequality $\phi^{(i)}(x_0) \leq 0$ also holds.

□

Therefore, if $2^n \leq |i| < 2^{n+1}$, $\phi^{(i)}(x_0) \leq \phi^{(i)}_{n+1}(x_0) = -|i|(3/4)^{n+1} \leq -2^n \cdot (3/4)^{n+1} = -3/4 \cdot (3/2)^n$. Finally, $\sum_{i\in\mathbb{Z}} \exp(\phi^{(i)}(x_0)) \leq 1 + \sum_{n=0}^{\infty} 2^{n+1} \exp(-3/4 \cdot (3/2)^n) = M < \infty$.

3.3. Construction step II. We will execute the same argument for the Cantor set C instead of the single point x_0. Since $R^{-2^n}C, \dots, C, \dots, R^{2^n-1}C$ are disjoint compact sets, there exists an ε-neighbourhood N of C such that $R^{-2^n}N, \dots, N, \dots, R^{2^n-1}N$ are disjoint. Define a bump function f so that $\operatorname{supp} f \subset N$, $f(x) = (3/4)^n$ on C and $0 \le f(x) < (3/4)^n$ on $N \setminus C$. Now we apply the same argument as in the previous subsection,to obtain the function $\phi \in C(S^1)$ such that $\sum_{i\in\mathbb{Z}} \exp(\phi^{(i)}(x)) < M < \infty$ for any $x \in C$.

We define a finite measure $\tilde{\mu}$ on S^1 by

$$\tilde{\mu} := \sum_{i\in\mathbb{Z}} (\exp \circ \phi^{(i)} \circ R^{-i}) R^i_* \mu_0.$$

Normalize $\tilde{\mu}$ to obtain a probability measure μ, namely $\mu := \frac{\tilde{\mu}}{\int_{S^1} d\tilde{\mu}}$.

Define h and F as in the section 2, then

$$\frac{dF^{-1}_* \, Leb}{d \, Leb} = \frac{dR^{-1}_* \mu}{d\mu} \circ h = g \circ h$$

is a continuous function because $g(x) = \exp(\phi(x))$. We proved Theorem 1.1 for the case $\alpha = (\sqrt{5} - 1)/2$.

4. Construction of Cantor set for general α

For general irrational number α, it is enough to find a Cantor set C that satisfies Claim 2.1. For a real number x and a function $p \colon \mathbb{N} \to \mathbb{N}$, define the approximation constant $c_p(x)$ by

$$c_p(x) := \liminf_{q\to\infty} \left(p(q) \cdot \operatorname{dist}(x, \tfrac{1}{q}\mathbb{Z}) \right).$$

A real number x is said to be p-approximable if $c_p(x) = 0$. Note that x is well approximable if x is p-approximable for $p(q) = q^2$, so this is a generalization of well-approximability.

It is clear that $c_p(x) = 0$ if x is rational number. On the other hand, for any irrational number x we can find a function p satisfying $c_p(x) > 0$. Moreover, we will show the following lemma.

Lemma 4.1. *For a given irrational number α, we can find a function p such that $c_p(m\alpha) \ge 1$ for any nonzero integer m.*

Proof. Since $c_p(-mx) = c_p(mx)$, it is enough to show the lemma for the case $m \in \mathbb{N}$. Let us start for any natural numbers n and q by taking a natural number $p_n(q)$ so that $p_n(q) \cdot \operatorname{dist}(n\alpha, \frac{1}{q}\mathbb{Z}) \ge 1$. Then define a function p by

$$p(q) = \max_{1\le n\le q} p_n(q).$$

By this construction $p(q) \ge p_m(q)$ for any $q \ge m$, therefore

$$\begin{aligned} c_p(m\alpha) &= \liminf_{q\to\infty} \left(p(q) \cdot \operatorname{dist}(m\alpha, \tfrac{1}{q}\mathbb{Z}) \right) \\ &\ge \liminf_{q\to\infty} \left(p_m(q) \cdot \operatorname{dist}(m\alpha, \tfrac{1}{q}\mathbb{Z}) \right) \\ &\ge 1. \end{aligned}$$

□

For this function p, we inductively take an increasing sequence $q_0, q_1, \dots$ of natural numbers satisfying the following conditions; $q_0 = 1$, $q_n | q_{n+1}$, $q_n/q_{n+1} \leq 1/3$ and $p(q_n)/q_{n+1} \leq 2^{-n}$. Define a Cantor set C by

$$C := \Big\{ \sum_{n=1}^{\infty} \frac{\varepsilon_n}{q_n} \mid \varepsilon_n = 0 \text{ or } 1 \Big\}.$$

This Cantor set C consists of p-approximable numbers. We can also show the following lemma.

Lemma 4.2. *For any $\beta \in C - C$, the approximation constant $c_p(\beta)$ is equal to 0, where*

$$C - C = \Big\{ \sum_{n=1}^{\infty} \frac{\varepsilon'_n}{q_n} \mid \varepsilon'_n = 0 \text{ or } \pm 1 \Big\}.$$

Proof.

$$\begin{aligned} p(q_i) \cdot \operatorname{dist}(\beta, \tfrac{1}{q_i}\mathbb{Z}) &\leq p(q_i) \left| \sum_{n=i+1}^{\infty} \frac{\varepsilon'_n}{q_n} \right| \\ &\leq p(q_i) \sum_{n=i+1}^{\infty} \frac{1}{q_n} = \frac{p(q_i)}{q_{i+1}} \sum_{n=i+1}^{\infty} \frac{q_{i+1}}{q_n} \leq \frac{1}{2^i} \sum_{k=0}^{\infty} \left(\frac{1}{3}\right)^k = \frac{3}{2^{i+1}}. \end{aligned}$$

Thus

$$\begin{aligned} c_p(\beta) &= \liminf_{q \to \infty} \Big(p(q) \cdot \operatorname{dist}(\beta, \tfrac{1}{q}\mathbb{Z}) \Big) \\ &\leq \liminf_{i \to \infty} \Big(p(q_i) \cdot \operatorname{dist}(\beta, \tfrac{1}{q_i}\mathbb{Z}) \Big) \\ &\leq \liminf_{i \to \infty} \frac{3}{2^{i+1}} \\ &= 0. \end{aligned}$$

Therefore $c_p(\beta) = 0$. □

Claim 2.1 follows from Lemma 4.1 and Lemma 4.2, so we proved Theorem 1.1 for the general case.

References

[D] A. Denjoy, *Sur les courbes défini par les équations différentielle à la surfase du tore.* J. Math. Pures Appl. 9(11) (1932), 333-375.

[H] M. R. Herman, *Sur la conjugaison différentiable des difféomorphismes du cercle a des rotations.* Publ. Math. I. H. E. S., 49(1979), 5-242.

[KH] A. Katok and B. Hasselblatt, *Introduction to the Modern Theory of Dynamical Systems.* Encyclopedia of Mathematics and its Applications, Vol. 54, Cambridge University Press, 1995.

[OR] F. Oliveira and L. F. C. da Rocha, *Minimal non-ergodic C^1-diffeomorphisms of the circle.* Erg. Th. Dyn. Sys. 21(2001), 1843-1854.

Graduate School of Mathematical Sciences, The University of Tokyo, 3-8-1 Komaba, Meguro-ku, Tokyo, 153-8914 Japan

E-mail address: `kodama@ms.u-tokyo.ac.jp`

Department of Mathematics, College of Science and Technology, Nihon University, 1-8-14 Kanda, Surugadai, Chiyoda-ku, Tokyo, 101-8308 Japan

E-mail address: `matsumo@math.cst.nihon-u.ac.jp`